


PERLA



No. 4 December 1979

BRIGHAM YOUNG UNIVERSITY

NOTICES

The Seventh International Symposium on Plecoptera will be held August 19-22, 1980, at the Nara Women's University in Nara, Japan. See page II for details.

The Plecoptera meetings are being held in conjunction with the XVI International Congress of Entomology (Kyoto, 3-9 August, 1980) and the XXI Congress of the International Association of Theoretical and Applied Limnology SIL (Kyoto, 24-31 August, 1980).

PERLA

A Newsletter for Plecopteroologists

EDITORS:

**Richard W. Baumann, Monte L. Bean Life Science Museum,
Brigham Young University, Provo, Utah 84602**

**Peter Zwick, Limnologische Flussstation, Max-Planck-
Instituts für Limnologie, Postfach 102, BRD 6407,
Schlitz, West Germany**

EDITORIAL ASSISTANTS:

**Carol Cutler
Jean Stanger**

Notes on a new classification of Plecoptera

Late in 1977, shortly before PERLA 3 was published, Prof. KEVAN of McGill University, Ste. Anne de Bellevue, Quebec, sent a copy of his "Suprafamilial classification of 'orthopteroid' and related insects" (KEVAN 1976, 1977a, b). This includes a classification of Plecoptera. Permission to reproduce this with some notes in PERLA was readily granted (for which Prof. KEVAN is sincerely thanked), but it arrived shortly after PERLA 3 had been published. This information is therefore presented now after considerable delay but will probably be new for many of us. My brief account is inevitably biased by my own views and is intended to direct those interested to the original paper. Prof. KEVAN informed me that reprints are no longer available, but the text forms part of Memoir 4 of the Lyman Entomological Museum and Research Laboratory, McGill University, and can be obtained for \$0.00/copy.

The present higher classification of orthopteroid insects and many other groups is described as being in a state near anarchy, if not chaos, or as a fluid situation at least. The author is concerned about numerous conflicting and frequently changing classifications and about the way in which the non-specialist is left behind with outmoded classifications differing from one text to the other. He is also alarmed about the situation of those who have to learn or even to teach insect taxonomy. No doubt these are serious problems calling for a change.

Rational action by entomologists and zoologists in general (extension of the scheme to all animal classification is proposed) is required. KEVAN proposes to start putting our house in order by first deciding where to put the principal furniture and by trimming untidy pieces later. We should, unlike Alice in Wonderland, first decide on appropriate sizes for slices, then cut up cakes and distribute them. If furniture or cakes were concerned, I would have no objections. However, these examples have little in common with our situation. The cakes we are dealing with were baked by evolution long ago and in many cases with time have lost shape to the extent that they have become unrecognizable. We do not have to cut up cakes, but we have to reconstruct them from crumbs. Our task is also not to

arrange existing furniture but to make it from bits and pieces that we have to find, recognize and fit together. In doing this we must constantly check to see if a screw we may find in some untidy corner indicates that some of the furniture we have already assembled was put together incorrectly.

The major problems of stonefly classification arose not because specialists dealt with untidy pieces and forgot about the principal furniture, but because students of Plecoptera were so sure of what their principal furniture was. The higher classification of Plecoptera can readily be traced back to LATREILLE. Unfortunately, Filipalpia, on which plecopterologists relied on so long was not their house but a pack containing a refrigerator and washing machine.

Of course, KEVAN allows for modifications of classifications if need be. "Here the phyletic or cladistic approach to taxonomy will play a dominant role. The result could then be put to a test by the use of 'phenetic' methods. Indeed, I believe that it is here that 'numerical taxonomy' has its most important role to play, particularly in respect of ranking the taxa recognized within a given hierarchical system..." But is it inevitable that such modifications bring about nomenclatorial changes upsetting textbooks and puzzling non-specialists?

KEVAN proposes a scheme which "aims at minimal disturbance of existing practice, and even where details may be unfamiliar, it is seldom actually innovative." An existing...hierarchical system is adopted uniformly... The principles of symbolic logic (GREGG, 1954) are employed, so that all superior categories are divided into an equal number of inferior categories of equivalent rank. The number of subcategories used is the optimum workable on the basis of maximum need within any group. To achieve this, each 'primary' category falls within a 'super-'category and includes within it one or more 'sub-' and 'infra-'categories." The rank of every taxon is indicated by a special suffix like those prescribed for family-group names by ICZN or like those generally used by botanists. A change in rank is made by change of the suffix, but author and date are not changed. They

may be put into parentheses to indicate some later change of form and/or rank, and author and date of such changes may be added. It is very important that absolute ranking to which much weight is assigned in this scheme "for the present... must largely be subjective" (KEVAN 1977b).

Stability of the scheme arises from extension of the principle of ICBN "for family-group names and their 'co-ordination' up to and including the level of suborder. The same rule of priority is applied. (Priority is also applied, in a modified way, to the names of orders and higher taxa but these are not coordinate with family-group names)." Extension of priority to before LINNAEUS, 1758, is requested for names above sub-order, and priority goes first to a name with a standard termination (see below). "Whether or not an author used a name in a restricted or expanded sense ... is of no consequence in the proposed system." This is so, because the names are considered *per se*, separate from their sense and the concepts they were used for. The point of reference is the name of the genus from which the oldest family-group name was derived.

Plecoptera are sometimes thought to be related to Orthoptera (though KEVAN is not convinced they are) and have therefore been included in the scheme. We can therefore test it on familiar grounds. The following is the classification proposed by KEVAN for the placement of Plecoptera, and their subdivision. In the first section, I have usually omitted synonyms, but the classification of Perlariae is copied in detail; abbreviations and symbols have been replaced by full wording, and an accidentally misspelled name has been corrected.

Superphylum: INSECTA Linnaeus, 1758 (after Plinius, 77 A.D., below); ARTICULATA Cuvier, 1805 (part); CONDYLOPA (Condylopes) Latreille, 1831 ARTHIROPODA Siebold + Stannius, 1845 or EUARTHIROPODA auctt. (polyphyletic).

Phylum: ENTOMA (Aristoteles, ca. 330 B.C., part) Latreille 1796 (part);...

Subphylum: TRACHEATA Haeckel, 1866;...

Infra phylum: ATELOCERATA Heymons, 1901, 1909;...

- Superclass:** HEXAPODA Latreille, 1825;...
- Class:** INSECTA (*Caius Plinius Secundus*, 77 A.A., part)
Linnaeus, 1758 (part; ectognathous hexapods only),
sensu Hennig, 1956;...
- Subclass:** DICONDYLIA Hennig, 1953.
- Infraclass:** PTILOTA (Aristoteles,...winged "Entoma" ...)
- Supercohort:** NEOPTERYGOTA Crampton, 1924, 1926, 1928, 1929;... (KEVAN proposes use of this junior synonym, because TETRAPTERA Aristoteles and several others, which have priority, are "too general and/or ambiguous" to be used)
- Cohort:** POLYNEPHRIA Brauer, 1885 (part)...
(if procedures proposed by KEVAN are to be followed, the oldest synonym, AMETABOLA Burmeister, 1829, should be used, because it is not a homonym of AMETABOLIA Leach, 1815)
- Subcohort:** SYNISTATA Fabricius, sensu restricto...
- Infracohort:** PERLARIAE Latreille, 1802; PERLARIDES Leach, 1815;
NEMOURACDES Billberg, 1820; PERLARIA Newman, 1834.
(the other infracohorts of Synistata are PROTOPERLAFIAE Tillyard, 1928, s.l., and EMBIARIAE Handlirsch, 1903)
- Superorder:** PLECOPTEROIDA (Burmeister) Kevan, 1976.
- Order:** Plecoptera Burmeister, 1800.
- Suborder:** Eustenolodes (Tillyard) Kevan, 1976; Archiperlaria lilles, 1965 (with additions); Antarctoperlaria Zwick, 1973.
- Infraorder:** Eustheniidea (Tillyard) Kevan, 1976.
- Superfamily:** Eustenoloides (Tillyard) Kevan, 1976 ---idae Tillyard, 1921.

- Infraorder:** *Gripopterygidea* (Enderlein) Kevan, 1976.
- Superfamily:** *Austroperloidea* (Tillyard) Illies, 1965 (part)-- -idae Tillyard, 1921.
- Superfamily:** *Gripopterygoidea* (Enderlein) Kevan, 1976-- -idae Enderlein, 1909.
- Suborder:** *Nemourodea* (Billberg) Kevan, 1976; *Nemourina* Albards, 1889; *Filipalpia* Klapalek, 1905; *Holognatha* Enderlein, 1909; *Arctoperlaris* Zwick, 1973 (part).
- Infraorder:** *Nemouridae* (Billberg) Kevan, 1976.
- Superfamily:** *Taeniopterygoidea* (Enderlein) Illies, 1965 (part)-- -idae Enderlein, 1909.
- Superfamily:** *Nemouroidea* (Billberg) Bradley, 1946-- *Nemouraedes* Billberg, 1820 (part) (-idae Selys, 1888).
- Suborder:** *Perloidea* (Latreille) Bradley, 1946 (as superordinal name); *Subulipalpia* Klapalek, 1905; *Setipalpia* Enderlein, 1909 (part); *Systellognatha* Enderlein, 1909; *Arctoperlaris* Zwick, 1973 (part).
- Infraorder:** *Pteronarcidae* (Yakobson + Blanki) Kevan, 1976.
- Superfamily:** *Pteronarcoidea* (Yakobson + Blanki) Kevan, 1976-- -int Yakobson + Blanki, 1904 (-idae Enderlein, 1909).
- Infraorder:** *Perlidae* (Latreille) Karach, 1883.
- Superfamily:** *Perloidea* (Latreille) Mandlirach, 1903 -- -idae Latreille, 1802 (~ides Leach, 1817; -idae Stephenson, 1836).

The second superorder within PERLARIAE is MIONOPTEROIDA (Martynov, part) Kevan, 1976.

As explained before, names are considered per se; references are to the first use of the names, but must not indicate that the author's concept is accepted. Hence, it is no contradiction if, for example, Holognatha Enderlein (which included all Eustheniodea), and Filipalpia Klapalek (which included Pteronarcidae) are listed as synonyms of the much more restricted Nemouridea Billberg. All included and were, so to speak, based on Nemoura, the type genus of the oldest family-group name. An extension of the principle of coordinate categories to subordinal names would in fact make them synonyms. (Setipalpia Enderlein 1909 might be an error for Setipalpia Klapalek, 1909, because Enderlein refused to use Setipalpia and proposed Systalognatha instead). There is no example in Plecoptera, but a group name derived from some type genus would replace names formed otherwise (e.g., Setipalpia, Holognatha, Arctoperlaria), even if these had priority.

This flexibility in the use of names also means that, unless listed, it is not possible to say which families are included in a given group except the one containing the genus from which the group name was derived. Position of Diamphipnoidae, Scopuridae, Capniidae, Leuctridae, Notonemouridae, Peltoperlidae, Perlodidae and Chloroperlidae cannot be read from the scheme, nor can it be inferred reliably from the references given. It is of course easy to guess where they would probably be placed, because the scheme is obviously strongly influenced by ILLIES (1965). However, such vagueness might be important as a requirement for the availability of names, provided ICZN would be changed as proposed by KEVAN.

The system I have proposed (1969, 1973) included fundamental changes of systematic concepts and of names. If KEVAN's proposals were followed, the nomenclatorial consequences would be much less drastic. As he rightly explained in a letter, his scheme would describe my classification like this (in an abbreviated form):

Order: Plecoptera Burmeister

- Suborder: Eustheniodea Tillyard (*Antarctoperlaria* Zwick)
- Infraorder: Eusthenidea Tillyard
- Infraorder: Gripopterygidea Enderlein
- Suborder: Perlodea Latreille (*Nemouridea* Billberg;
Arctoperlaria Zwick)

Infraorder: Nemouridea Billberg
Infraorder: Pteronarcides Yokobson + Bianki
Infraorder: Perlidea Latreille

Such stability of nomenclature is really impressive. In the present state of knowledge, it cannot be stable if it reflects our concepts. To obtain stability, the principle of coordinate categories is used to strip names of their sense and then a stable list of names can be drawn. This, to me, appears rather senseless. If such restrictions were accepted, taxonomic texts would really not often have to change. Hopefully, their users will not be satisfied by learning lists of empty names. Rather, they will remain interested to know that (as I put it before) plecopteroologists have been living on their refrigerator and washing machine for 150 years and will be interested to know why this happened. Those who have to (but sometimes do not like to) pay insect taxonomists, will be pleased to have it documented that taxonomists have achieved near to nothing since LATREILLE and BILLBERG, and that systematic problems have been solved, if not by LINNAEUS, then not long after him.

Of course I realize that the present ruling (as proposed by KEVAN) is already established for taxa up to superfamilies. This is the realm of almost innumerable taxa, the domain of specialists with a knowledge of problems and concepts, be they apparent in nomenclature or not. The number of taxa to which KEVAN wants the present ruling extended is relatively modest, and this is the area of interest to non-specialists. They need help, even though taxonomy must always be expected to move as long as it is a living science.

I believe that help is under way, but from the opposite direction than the one suggested by KEVAN. "Let us not worry, therefore, about stability as it applies to biological concepts in classification; rather, we should be concerned about ways of achieving better phylogenetic analyses. Only as phylogenetic analyses progress toward an evergreater degree of probability in expressing the actual path of evolution will our classifications become increasingly stable" (The late H. D. ROSS in a discussion; reprint without bibliographical data). Progress could be made almost accidentally only as long as no scientific theory of systematics was available. However, almost 200 years after LINNAEUS, it has become available through the works of HENNIG. Certainly, the English translation of his book on insect

phylogeny (which is presently being prepared) will provide strong stimuli. I also agree with GRIFFITHS (1976) that we should free ourselves of Linnean categories. These categories had been proposed for what was thought to be classification of an unchanging product of creation, but we try to apply the same categories in the reconstruction of the evolutionary process, which LINNAEUS did not understand. His categories were meant as a support, but on different, theoretical grounds they have turned into a fence. We should leap over this instead of setting up new and even higher fences. Instead of extending this practice, ICZN should stop prescribing suffixes for categorical levels. The scale of categories is rigid and logical but it has nothing to do with evolution.

In a letter, Prof. KEVAN said that giving up fixed categorical levels would be retrograde and would lead to anarchy (which we are said to be close to anyway). This need not be. Nomenclature was correctly called our book-keeping by H. H. ROSS and modern technology should allow us to fit our book-keeping to our results, and no longer do the reverse.

There seems to be no need for a full bibliography here, so I list only the two papers describing the controversial views most clearly. Prof. KEVAN's paper was first presented, and a limited first edition of copies was distributed, during the XV International Congress of Entomology, Washington (1976). A second edition appeared as 'Notes from the Lyman Entomological Museum and Research Laboratory, No. 2' (1977a) and the complete text (1977b) is the one listed below.

GRIFFITHS, D.C.G. (1976): The Future of Linnean Nomenclature. - Syst. Zool., 25: 168-173.

KEVAN, D.K.McE. (1977): The Higher Classification of the Orthopteroid Insects: a General View. (Papers presented at Section 1 Symposium, XV International Congress of Entomology, Washington, D.C., U.S.A., August, 1976). - Memoirs of the Lyman Entomol. Mus. and Res. Lab., 4:1-31, plus Appendix of 26 pages.

Peter ZWICK
Limnologische Flussstation
der MPG, Postfach 260
D-6407 Schlitz

Seventh International Symposium on Plecoptera

At the Sixth International Symposium on Plecoptera in Schlitz, West Germany, in 1977, it was decided that the seventh meeting should be held in Nara, Japan. Dr. Teizi Kawai of the Nara Women's University is the organizer for this event. The symposium will be held August 19-22, 1980 at University Hall, Nara Women's University, Nara, Japan.

In addition to the presentation of papers, a field trip to Yoshino will take place. We will spend some time there at a typical mountain stream, the Talamigawa River, which is a branch of the Yoshino River. The field trip will conclude a day before the opening of the XXI SLL Congress in Kyoto.

Nara is a famous ancient city in the Kinki District of Japan. It is located in a rich fertile area in which Japanese culture has flourished from earliest times. Since the Asuka period of 552 to 646, Nara has been a region in which the finest monuments of Japanese creative genius and the superb natural beauties of the area have blended into a surpassing harmony. It will be a beautiful setting for our meetings.

For those of you who wish to present a paper, please inform Dr. Kawai of the title, time requirements, and any special equipment that you might need. A standard size 5 cm x 5 cm slide projector, a 16 mm movie projector, and a tape recorder will be available. Please also indicate any accompanying persons. The fee for the symposium will be 15,000 yen (60 US dollars) for each person.

At the conclusion of the conference, a bulletin of the proceedings of the symposium, including abstracts from the papers presented, as well as the names and addresses of those who attended will be published. Please notify Dr. Kawai soon if you plan to attend. The registration deadline is January 31, 1980. We look forward to seeing you there.

continued page 12

For more information please write to:

**Dr. Teizi Kawai
Biological Institute
Nara Women's University
Kitauoya - Nishi - Machi
630 Nara, Japan**

Bulletin Board

NEWS: Ken Stewart and Bill Stark are continuing their work on a book of the nymphs of North American stoneflies. They are being funded by National Science Foundation. We encourage all our North American colleagues to support this study which will be of benefit to everyone interested in aquatic insects.

New Journal

A new journal entitled AQUATIC INSECTS is being published on a quarterly basis by Swets Publishing Service. Prof. Dr. J. Illies is the editor-in-chief for this new periodical. AQUATIC INSECTS includes both taxonomy and ecology of aquatic insects, as well as papers on aquatic insects that were previously widely scattered in various entomological and limnological journals. Orders can be placed via a subscription agent or direct with the publisher. Subscriptions must only be made on a calendar year basis. The price for individuals is 93.50 Dutch guilders (US \$45.60), and 128.50 Dutch guilders (US \$62.75) for institutions. Customers in North America should write to:

Swets North America, Inc.
P. O. Box 517
Bala Cynwyd, PA 19312
USA

Customers in other countries should order through:

Swets Publishing Service
P. O. Box 825
2160 SZ Lisse
The Netherlands

A. R. Gaufin Stonefly Collection

The stonefly collection which Arden R. Gaufin and students have assembled over the past 30 years has been placed on loan for a 5 year period beginning 1980. The collection is now at the Monte L. Bean Life Science Museum, Brigham Young University, Provo, Utah 84602. Hopefully at the end of the 5 year loan period the collection will find a permanent home in Provo, Utah.

All inquiries and loan requests concerning the Gaufin collection should be sent to Richard W. Baumann at the above address.


Jewett Donates Stoneflies to Institutions

We have been informed that Stanley G. Jewett, Jr. has recently donated his collection of domestic aquatic insects to Oregon State University and his exotic aquatic insects to the California Academy of Sciences.

**California Academy of Sciences
Golden Gate Park
San Francisco, CA 94118**

**Oregon State University
Department of Entomology
Corvallis, OR 97331**

Recent Plecoptera Literature



This section includes the Plecoptera papers published since Perla 3 was mailed two years ago. In the future the plan is to publish Perla every two years and include a literature section in every issue.

Please help us to make this section as complete and correct as possible by sending us copies of your publications and/or notes on errors found.

- AHSAN, M. (1974). Aquatic fauna of Swat Valley, Pakistan; Part II. A preliminary report on freshwater insects. *Biologia (Lahore)*. 20: 173-178. (In English).
- ALIMOV, A. F. & N. V. SHADRIN. (1977). Chloric content of some representatives of freshwater benthos. *Hydrobiol. Zh.* 13: 80-86. (Russ. with Engl. summ.).
- ALLAN, J. D. (1978). Trout predation and the size composition of stream drift. *Limnol. Oceanogr.* 23: 1231-1237.
- ANDERSEN, T. A., FJELLHEIM, R., LARSEN & C. OTTO. (1978). Relative abundance and flight periods of Ephemeroptera, Plecoptera and Trichoptera in a regulated West Norwegian river. *Norw. J. Ent.* 25: 139-144.
- ANDERSON, N. H. & J. R. SEDELL. (1979). Detritus processing by macroinvertebrates in stream ecosystems. *Annual Reviews of Entomol.* 24: 351-377.
- ARMITAGE, P. D. (1978). Downstream changes in the composition, numbers and biomasses of bottom fauna in the Tees below Cow Green Reservoir and in an unregulated tributary Maize Beck, in the first five years after impoundment. *Hydrobiologia* 58: 145-156.
- ASHLEY, D. L., D. C. TARTER & W. D. WATKINS. (1976). Life history and ecology of Diploperla robusta Stark and Gaufin (Plecoptera: Perlodidae). *Psyche* 83: 310-318.
- BAGGE, P. & V. SALMELA. (1978). The macrobenthos of the River Tourujoki and its tributaries (Central Finland): I. Plecoptera, Ephemeroptera and Trichoptera. *Not. Entomol.* 58: 159-168.
- BALDUZZI, A., E. GRAINO & S. SPANO. (1976). The Arroscia-Centa hydrologic system (western Liguria) in relation to its zoobenthonic populations, with special attention to Ephemeroptera and Plecoptera. *Boll. Mus. Inst. Biol. Univ. Genova* 44: 25-54. (Ital. with Engl. summ.).

- BAUMANN, R. W. (1976). An annotated review of the systematics of North American stoneflies (Plecoptera). *Perla* 2: 21-23.
- BAUMANN, R. W. (1977). Water insects and their relatives. *Amer. Bio. Teacher* 39: 295-299.
- BAUMANN, R. W. (1979a). Stoneflies of the Rocky Mountains. *Gewasser und Abwasser* 64: 48.
- BAUMANN, R. W. (1979b). Nearctic stonefly genera as indicators of ecological parameters (Plecoptera: Insecta). *Gt Basin Nat.* 39: 241-244.
- BAUMANN, R. W., A. R. GAUFIN & R. F. SURDICK. (1977). The stoneflies (Plecoptera) of the Rocky Mountains. *Memoirs Amer. Entomol. Soc.* 31, 208 pp.
- BEDNARIK, A. F. & W. P. McCAFFERTY. (1977). A checklist of the stoneflies or Plecoptera of Indiana. *Great Lakes Entomol.* 10: 223-226.
- DENFIELD, E. F., D. S. JONES & M. F. PATTERSON. (1977). Leaf pack processing in a pastureland stream. *Oikos* (Copenhagen) 29: 99-103.
- BRNGTSSON, J. (1977). Food preference experiments with nymphs of Nemoura cinerea (Retz.) (Plecoptera). *Flora Fauna* 83: 36-39. (Ger. with Engl. summ.).
- BERTHELEMY, C. (1979). Mating calls and taxonomy in Pyrenean Isoperla. *Gewasser und Abwasser* 64: 71-72.
- BLAHAK, P. (1970). Contribution to the knowledge of the relationship of zoobenthos to food of brook trout and of grayling. *Acta rer. natur. Mus. Nat. Slov. (Bratislava)* 24: 41-81. (Czech. with German summ.).
- BLOESCH, J. (1977). Bodenfaunistische Untersuchungen in Aare und Rhein Schweizer. *Z. Hydrobiol. (Basel)* 39: 46-68.

- BONAZZI, G. & P. F. GHETTI. (1977). The macroinvertebrate populations of Parma Stream: Results of a yearly cycle of research. Ateneo. Parmense Acta. Nat. 13: 351-396. (Ital. with Engl. summ.).
- BRAASCH, D. & W. JOOST. (1977). *Leuctra kumanekii* n. sp. --eine neue spätere Steinfliege (Plecoptera, Leuctridae) aus Bulgarien. Ent. Nachr. 21: 183-185.
- BRAASCH, D. & E. ODRICH. (1975). *Amphinemura borealis* (Morton 1894) - a new plecopteran for the DDR. Ent. Nachr. 19: 68-70. (Ger. with Engl. summ.).
- BRITTAINE, J. E. (1977). The effect of temperature on the egg incubation period of *Taeniopteryx nebulosa* (Plecoptera). Oikos (Copenhagen) 29: 302-305.
- BRITTAINE, J. E. (1978). Semivoltinism in mountain populations of *Nemurella pictetii* (Plecoptera). Oikos (Copenhagen) 30: 1-6.
- BRITTAINE, J. E. (1979). Semivoltinism in mountain populations of *Nemurella pictetii* (Plecoptera). Gewässer und Abwasser 64: 29 (Abs.).
- BRITTAINE, J. E. & A. LILLEHAMMER. (1978). The fauna of the exposed zone of Øvre Heimdalvatn: methods, sampling stations and general results. Holarct. Ecol. (Copenhagen) 1: 221-228.
- BRODSKIY, A. K. (1979). Evolution of the flight apparatus in Plecoptera. Part 1. Functional morphology of wings. Entomol. Obozr. (Leningrad) 58: 69-77. (In Russian).
- BROOKER, M. P. & R. J. HEMSWORTH. (1978). The effect of the release of an artificial discharge of water on invertebrate drift in the R. Wye, Wales. Hydrobiologia (The Hague) 39: 155-163.
- CASPERS, N. & H. STIERS. (1977). Contribution to the knowledge of the Plecoptera of the Eifel (Insecta: Plecoptera). Decheniana 130: 136-150. (Ger. with Engl. summ.).
- CHINO, Y. (1978). The female of *Nogiperla uenoii* (KOHNO) (Plecoptera, Peltoperlidae). New Entomol. 27: 9-17.

- CLIFFORD, H. F. (1978). Descriptive phenology and seasonality of a Canadian brown-water stream. *Hydrobiologia* (Den Haag) 58: 213-231.
- COIFFAIT, H. (1976). Recent works on taxonomy and biogeography. *Nouv. Rev. Ent.* 6: 325-332. (In French) (Plecoptera references).
- CONSIGLIO, C. (1978). On some stoneflies of the genus Neoperla from Ethiopia (Plecoptera). *Accad. Nazion. dei Lincei* 243: 59-61.
- COWIE, D., A. J. CONNER & L. N. CONNER. (1978). A survey of the benthic invertebrates from the Fresh Water Valley, Stewart Island, New Zealand. *Mauri Ora* 6: 27-32.
- CRISP, D. T., R. H. K. MANN & J. C. MCCORMACK. (1978). The effects of impoundment and regulation upon the stomach contents of fish at Cow Green, Upper Teesdale. *J. Fish. Biol.* 12: 287-302.
- CUSIMAN, R. M., J. W. ELWOOD & S. G. HILDEBRAND. (1975). Production dynamics of Alloperla mediana Banks (Plecoptera: Chloroperlidae) and Diplectrona modesta Banks (Trichoptera: Hydropsychidae) in Walker Branch, Tennessee. Oak Ridge Natl. Lab. Environ. Sci. Div. Publ. 785.
- CUSIMAN, R. M., J. W. ELWOOD & S. G. HILDEBRAND. (1977). Life history and production dynamics of Alloperla mediana and Diplectrona modesta in Walker Branch, Tennessee. *Am. Mid. Nat.* 98: 354-364.
- DAUTI, E. (1977). Contribution to the study of the fauna of Plecoptera in the river system of upper flow of the Nerodimka River. *Acta Biol. Med. Exp.* 2: 65-70.
- DEVONPORT, B. F. & M. J. WINTERBOURN. (1976). The feeding relationships of two invertebrate predators in a New Zealand river. *Freshwat. Biol.* (Oxford) 6: 167-176.
- DIEM, F. (1977). Über die Fauna des Niklbaches (Naturschutzgebiet Sengsengebirge O. O.) mit besonderer Berücksichtigung der Steinfliegen (Insecta: Plecoptera). Hausarbeit Lehramtsstudium Universität Innsbruck; unpublished manuscript, 111 pp.

- DONALD, D. B. & R. S. ANDERSON. (1977). Distribution of the stoneflies (Plecoptera) of the Waterton River Drainage, Alberta, Canada. Can. Wildlife Service. Syesis 10: 111-120.
- DORGELO, J. (1976). Further observations on drift in two brooks in Auvergne. Ann. Stn. Biol. Besse-en-Chandesse 10: 233-249.
- DOSDALL, L. (1976). Saskatchewan winter stoneflies. Blue Jay 34: 198-202.
- DOSDALL, L. & D. M. LEMKUHL. (1979). Stoneflies (Plecoptera) of Saskatchewan. Quaestiones Entom. 15: 3-116.
- DURRENFELDT, A. (1978). Untersuchungen zur Besiedlungsbiologie von Kalktuff - faunistische, ökologische und elektronenmikroskopische Befunde. Arch. Hydrobiol. Suppl. 54: 1-79.
- DUTHER, S. G. (1977). Neuropteroidea from the river bed of the As of Drenthe. Entomol. Ber. 37: 81-85. (Neth. with Engl. summ.).
- FEY, J. M. (1977a). Die Aufheizung eines Mittelgebirgsflusses und ihre Auswirkungen auf die Zoozonose - dargestellt an der Lenne (Sauerland). Arch. Hydrobiol. Suppl. 53, Monogr. Beitr.: 307-363.
- FEY, J. M. (1977b). Ein Beitrag zur Biologie der Lenne. Der Sauerländer Naturbeobachter. Veröffentl. Naturwiss Verein Ludenscheid 11: 1-19.
- FIANCE, S. B. (1977). The genera of Eastern North American Chloroperlidae (Plecoptera): key to larval stages. Psyche 84: 308-316.
- FIELDS, W. C., Jr. (1977). A new Megaleuctra from California (Plecoptera: Leuctridae). Par-Pac. Entomol. 53: 211-214.
- FINNI, G. R. & L. CHANDLER. (1977). Postdiapause instar discrimination and life history of the capniid stonefly Allocapnia granulata (Clasen) (Insecta: Plecoptera). Am. Mid. Nat. 98: 243-250.

- PINNI, G. R. & L. CHANDLER. (1979). The micro distribution of Allocapnia nainda (Plecoptera: Capniidae). J. Kans. Entomol. Soc. 52: 93-102.
- FLANNAGAN, J. F. (1978). The winter stoneflies Allocapnia granulata (Capniidae), Taeniopteryx nivalis and T. parvula (Taeniopterygidae) in southern Manitoba, Canada Ent. 110: 111-112.
- FREDEEN, F. J. H. & D. T. SPURR. (1978). Collecting semi-quantitative samples of black fly larvae (Diptera: Simuliidae) and other aquatic insects from large rivers with the aid of artificial substrates. Quaestiones Entom. 14: 411-431.
- FRIBERG, F., L. M. NILSSON, C. OTTO, P. SJOSTROM, B. W. SVENSSON, B. SVENSSON, AND S. ULFSTRAND. (1977). Diversity and environments of benthic invertebrate communities in south Swedish streams. Arch. Hydrobiol. 81: 129-154.
- FROELICH, C. G. (1979). The genus Eutactophlebia. Gewasser und Abwasser 64: 70-71.
- FULLER, R. L. & K. W. STEWART. (1979). Stonefly (Plecoptera) food habits and prey preference in the Dolores River, Colorado. Am. Mid. Nat. 101: 170-181.
- FULLINGTON, K. E. (1979). The nymphs of North American Taeniopteryx. Gewasser und Abwasser 64: 35-36.
- GILLESPIE, D. M., J. D. ELDRIDGE & C. K. THOMPSON. (1975). A kinetic model for static bioassay of insecticides (Pteronarcys californica used). Water Res. 9: 817-820.
- GIUDICELELLI, J. (1975). Analysis of endemism in the fauna from running waters in Corsica. Ecol. Mediterr. 1: 133-148.
- GODREY, P. J. (1977). Diversity as a measure of benthic macro-invertebrate community response to water pollution. Hydrobiologia (The Hague) 57: 111-122.

- GORE, J. A. (1978). A technique for predicting in-stream flow requirements of benthic macroinvertebrates. *Freshwat. Biol.* (Oxford) 8: 141-151.
- GRANT, P. R. & R. J. MACKAY. (1969). Ecological segregation of systematically related stream insects. *Can. J. Zool.* 47: 691-694.
- GRAY, L. J. & J. V. WARD. (1978). Food habits of stream benthos at sites of differing food availability. *Am. Midl. Nat.* 102: 157-167.
- HAGIWARA, K. (1977). On the biology of a stonefly, Eocapnia niviles (Veno) (Plecoptera, Capniidae), during the snow-season in Ashu, the northern suburb of Kyoto. *Kontyu* 45: 421-430. (Jpn. with Engl. summ.).
- HARPER, P. P. (1977). Capniidae, Leuctridae, and Perlidae (Plecoptera) from Nepal. *Oriental Insects* (Delhi) 11: 53-62.
- HARPER, P. P. (1979). Observations on the early instars of stoneflies (Plecoptera). *GWASSER und ABWASSER* 64: 18-28.
- HARPER, P. P. & R. F. KIRCHNER. (1978). A new stonefly from West Virginia (Plecoptera: Chloroperlidae). *Proc. Entomol. Soc. Wash.* 80: 403-406.
- HART, D. S. & M. A. BRUSVEN. (1976). Comparison of benthic insect communities in six small Idaho Batholith streams (includes Plecoptera). *Melanderia* 23: 1-18.
- HELENEN, W. (1976). List of the insect species recorded new to Finland during 1971-1975. *Not. Entomol.* 56: 109-119. (Ger. with Engl. summ.).
- HUGGINS, D. G. (1978). Kansas Plecoptera. *Trans. Kans. Acad. Sci.* 81: 153.
- HULDEN, I., M. MEINANDER, O. NYBOM & U. SILFVERBERG. (1977). Deletions from the Finnish fauna I. *Not. Entomol.* (Helsingfors) 57: 11-12.

- HYNES, H. B. N., N. K. KAUSHIK, N. A. LUCK, D. L. LUSH, Z. S. J. STOCKER, R. R. WALLACE & D. D. WILLIAMS. (1974). Benthos and allochthonous organic matter in streams. *J. Fish. Res. Bd. Can.* 31: 545-553.
- IKONOMOV, P. (1978). Plecopteres de la plaine de Skopje. *Ann. Fac. Biol. Univ. (Skopje)* 31: 5-18.
- ILLIES, J. (1978a). Comparative measurement of emergence in Breitenbach 1969-1976. *Arch. Hydrobiol. (Stuttgart)* 82: 432-440.
- ILLIES, J. (1978b). Plecoptera. In: ILLIES, J. (ed.): *Limnofauna Europaea*. - 2nd edn. (Stuttgart). XVII: 264-273.
- ILLIES, J. (1979). Population dynamics of Plecoptera from Breitenbach, Schlitz. *Gewässer und Abwasser* 64: 47.
- IVERSEN, T. M. (1978). Life cycle and growth of three species of Plecoptera in a Danish spring. *Ent. Meddr. (Copenhagen)* 46: 57-62.
- IVERSEN, T. M. & B. L. MADSEN. (1977). Allochthonous organic matter in streams. *Folia Limnol. Scand.* 17: 17-20.
- JOOST, W. (1977). *Nemoura monae* n. sp. -eine neue Steinfliege (Plecoptera: Nemouridae) aus dem Einzugsgebiet der Teberda (UdSSR, Westkaukasus). *Ent. Nachr. (Dresden)* 21: 27-31.
- JOOST, W. (1978). 3.3. Ordnung Plecoptern (Steinfliegen). In: *Wasserinsekten Kulturbund der DDR* (ed.) (Leipzig) 88 pp.
- KACANSKI, D. (1978). Plecoptera alvia reke Neretve (Plecoptera of the River Neretva drainage). *Codien. Biol. inst. Univ. Sarajevo* 31: 57-58.
- KACANSKI, D. (1979). Some characteristics of the Plecoptera fauna in Bosnia and Herzegovina (Yugoslavia). *Gewässer und Abwasser* 64: 49-55.

- KAISER, E. W. (1977). Short notations on Danish stoneflies (Plecoptera). *Flora Fauna* 83: 85-87. (Dan. with Engl. summ.).
- KAPOOR, N. N. (1978a). Effect of osmotic stress on the 'chloride cells' and ATP-ase activity in the tracheal gills of stonefly nymphs. (Insecta). *Amer. Zoologist* 18: 344.
- KAPOOR, N. N. (1978b). Effect of salinity on the osmoregulatory cells in the tracheal gills of the stonefly nymph, Paragnetina media (Plecoptera: Perlidae). *Can. J. Zool. (Ottawa)* 56: 2608-2613.
- KAPOOR, N. N. (1979a). The effect of osmotic stress on the survival and respiration of the stonefly nymph, Paragnetina media. *Gewasser und Abwasser* 64: 37-38.
- KAPOOR, N. N. (1979b). Osmotic regulation and salinity tolerance of the stonefly nymph Paragnetina media. *J. Insect Physiol.* 25: 17-20.
- KAPOOR, N. N. & K. ZACHARIAH. (1978). The internal morphology of sensilla on the abdominal gills of Thaumatoperla alpina (Plecoptera: Eustheniidae; sub-family: Thaumatoperlinae). *Can. J. Zool.* 56: 2194-2197.
- KAPOOR, N. N. & K. ZACHARIAH. (1979). The morphology of flower-like sensilla of Thaumatoperla alpina (Eustheniidae). *Gewasser und Abwasser* 64: 44.
- KASTER, J. L. & G. Z. JACOBI. (1978). Benthic macro-invertebrates of fluctuating reservoir. *Freshwat. Biol.* 8: 283-290.
- KAWAI, T. & Y. ISOBE. (1979). Some taxonomic relationships of the genera Schistoperla BANKS and Kiotina KIAPALEK. *Gewasser und Abwasser* 64: 74-76.
- KAWECKA, B., M. KOWNACKA & A. KOWNACKI. (1971). General characteristics of the biocoenosis in the streams of the Polish High Tatras. *Acta Hydrobiol. (Krakow)* 13: 465-476.

- KELLOGG, R. L. & R. V. VAULKLEY. (1976). Seasonal concentration of Dieldrin in water, Channel Catfish, and catfish-food organisms, Des Moines River, Iowa 1971-73. *Pestic. Monit. J.* 9: 186-194.
- KHALAF, G. & H. TACHET. (1977). The dynamics of macroinvertebrate colonization of artificial substrates in a stream. *Ann. Limnol.* 13: 169-190. (Fr. with Engl. summ.).
- KIRCHNER, R. F. (1978). Plecoptera records from West Virginia Ent. News. 89: 206.
- KLAUSNITZER, B., G. FRIESE, W. HEINICKE, W. JOOST & C. MULLER. (1978). Bedrohte Insektenarten in der Deutschen Demokratischen Republik I. Beitrag. *Entomol. Ber. (Berlin)* 1978: 81-87.
- KNIE, J. (1977). Okologische Untersuchung der Käferfauna von ausgewählten Fließgewässern des Rheinischen Schiefergebirges (Insecta: Coleoptera). (Includes Plecoptera). *Decheniana (Bonn)* 130: 151-221.
- KNIGHT, A. W. & C. A. SIEGFRIED. (1977). A teratomorphic stonefly (Plecoptera: Nemouridae). *Great Lakes Entomol. (East Lansing, Michigan)* 10: 165.
- KONDRATIEFF, B. C. & J. R. VOSHELL, JR. (1979). A new species of *Diploperla* (Plecoptera: Perlodidae) from Virginia. *Ann. Ent. Soc. Am.* 72: 451-453.
- KOPUNEN, S. (1977). Light trap catches of insects at Revo, northernmost Finland. *Nat. Entomol.* 57: 53-57.
- KOVALAK, W. P. (1978a). Diel changes in stream benthos density on stones and artificial substrates. *Hydrobiologia (Den Haag)* 58: 7-16.
- KOVALAK, W. P. (1978b). On the feeding habits of Phaeonophora capitata (Plecoptera: Perlidae). *Great Lakes Entomo.* 11: 45-50.

- KOWNACKI, A. (1977). Biocenosis of a high mountain stream under the influence of tourism. 4. The bottom fauna of the stream Rybi Potok (the High Tatra Mts.). *Acta Hydrobiol. (Krakow)* 19: 293-312. (Engl. with Polish summ.).
- KOWNACKA, M. & G. MARCREITER. (1978). Die Zoobenthos - Gesellschaften des Piburger Baches (Oetztal, Tirol). *Int. Revue ges. Hydrobiol.* 63: 213-232.
- KRNO, I. (1977). Stonefly Leuctra quadrimaculata Kis, 1963, a new species for the fauna of Czechoslovakia. *Biologia (Bratislava)* 32: 893-894. (Slovak. with Engl. summ.).
- KUUSELA, K. (1976). On the emergence biology of Taeniopteryx nebulosa (L.) (Plecoptera). *Ann. Entomol. Fenn. (Helsinki)* 42: 121-132.
- KUUSELA, K. & H. PULKKINEN. (1978). A simple trap for collecting newly emerged stoneflies (Plecoptera). *Oikos* 31: 323-325.
- LANGDON, T. J. & A. B. SCHLESINGER. (1978). Vertical distribution of macroinvertebrates in the Missouri River, U.S.A., near Fort Calhoun nuclear power station. *Proc. Nebr. Acad. Sci. Affil. Soc.* 88: 16.
- LAPCHIN, L. (1977a). Microrepartition des invertebres benthiques dans deux ruisseaux a Salmonides de Bretagne. *Ann. Hydrobiol. (Paris)* 8: 319-332.
- LAPCHIN, L. (1977b). Utilisation de l'analyse des correspondances pour l'étude de la répartition des invertebres benthiques dans deux ruisseaux a Salmonides de Bretagne. *Ann. Hydrobiol. (Paris)* 8: 333-354.
- LEVANODOVA, I. N., V. Ya. LEVANEDOV & E. A. MAKARCHENKO. (1977). Fauna of water invertebrate from the reserve "Kedrovaya Pad." In: Freshwater fauna of the reserve "Kedrovaya Pad" Proceedings of the Institute of Biology and Pedology 45: 3-43. (In Russian).

- LILLEHAMMER, A. (1978). The Plecoptera of Øvre Heimdalvatn. *Holarct. Ecol.* 1: 232-238.
- LILLEHAMMER, A. (1979). Stonefly populations at high altitudes in Norway. *Gewässer und Abwasser* 64: 48-49.
- LILLEHAMMER, A. & J. P. BRITAIN. (1978). The invertebrate fauna of the streams in Øvre Heimdalalen. *Holarct. Ecol.* 1: 271-276.
- LINDEGAARD, C. (1979). A survey of the invertebrate fauna, with special reference to Chironomidae (Diptera) in the rivers Laxa and Kraka, northern Iceland. *Oikos (Copenhagen)* 32: 281-288.
- LUEDTKE, R. J. & M. A. BRUSVEN & F. J. WATTS. (1976). Benthic insect community changes in relationship to in-stream alterations of a sediment-polluted stream (Plecoptera included). *Melanderia* 23: 21-39.
- MACAN, T. T. (1977a). A twenty-year study of the fauna in the vegetation of a moorland fish pond. *Arch. Hydrobiol.* 61: 1-24.
- MACAN, T. T. (1977b). The fauna in the vegetation of a moorland fish pond as revealed by different methods of collecting. *Hydrobiologia* 55: 3-16.
- MCCAFFERTY, W. P. (1978). Pre-management assessment of aquatic macroinvertebrates in a small sedimentary drainage area of the Maumee and Lake Erie Basin. *Great Lakes Entomol.* 11: 37-44.
- MCLELLAN, I. D. (1979). New Zealand terrestrial stoneflies and some ideas on speciation. *Gewässer und Abwasser* 64: 56-59.
- MADSEN, B. L. (1977). Studies of population movements in insects associated with streams. *Zool. Revy.* 29: 22-29. (Dan. with Engl. summ.).
- MALMQVIST, B., L. H. NILSSON & B. S. SVENSSON. (1978). Dynamics of detritus in a small stream in southern Sweden and its influence on the distribution of the bottom animal communities. *Oikos (Copenhagen)* 31: 3-16.

MAMAEV, B. M. (1972). Key to insects classified according to their larvae. Manual for teachers. Prosveshchenie Moskva 1-400.

MARLIER, G. (1977). Aquatic insects. Nat. Belg. 58: 97-110, 116-178. (Fr.).

MARLIER, G. (1979). A hydrobiological mission to the Seychelles. Nat. Belg. 60: 44-58. (Fr.).

MASTELLER, E. C. (1977). An aquatic emergence trap on a shale stream of Western Pennsylvania. Melshelmer Entomol. Series 23: 10-15.

MAUCH, E. (1976). Leitformender Saprobitat fur die biologische Gewasseranalyse. Cour. Forsch. - Inst. Senckenberg (Frankfurt) 21: XXVII + 797 pp.

MENDL, H. (1972). Plecoptera und Diptera, Nematocera (Tipulidae: Limoniinae) aus Ligurien Ein Beitrag zur Plecopteran und Dipteren fauna Italiens. Mitt. Naturwiss. Arbeitskreis, Kempten (Allgäu) 16: 53-58.

MENDL, H. & K. MULLER. (1978). The colonization cycle of Amphinemura standfussi (Insecta: Plecoptera) in the Abisko area, Sweden. Hydrobiologia 60: 109-112.

MULLER, K. & H. MENDL. (1978). Development of wings in Cappnia atra in the Abisko area, Sweden (Plecoptera). Entomol. Tidskr. 99: 111-112.

MULLER, K. & H. MENDL. (1979). The importance of a brackish water area for the stonefly colonization cycle in a coastal river. Oikos 33: 272-277.

NAGELL, B. (1979). Capacity to survive anoxia or low oxygen concentrations of Nemoura cinerea larvae (Plecoptera). Gewasser und Abwasser 64: 39.

NAGELL, B. & T. FAGERSTROM. (1978). Adaptations and resistance to anoxia in Cloeon dipterum (Ephemeroptera) and Nemoura cinerea (Plecoptera). Oikos 30: 95-99.

- NEDIC, D. (1975). Some aspects of the population density of bottom fauna of the rivers Liva and Komarnica and their most important tributaries (English summary). Glas. Repub. Zavoda Zast. Prir. Mux. Titogradu 8: 81-91.
- NELSON, C. H. (1977). Position of "Leuctra" divisa Hitchcock within the family Leuctridae (Plecoptera). Ent. News 88: 235-240.
- NEVES, R. J. (1978). Seasonal succession and diversity of stoneflies (Plecoptera) in Factory Brook, Massachusetts, U.S.A. J. N. Y. Ent. Soc. 86: 144-152.
- NEWELL, R. L. & G. W. MINSHALL. (1976). An annotated list of the aquatic insects of southeastern Idaho, Part I: Plecoptera. Ct Basin Nat. 36: 501-504.
- NIHMO, A. P. (1978). (Book Review) The stoneflies of the Rocky Mountains by R. W. Beumann, A. R. Gaafin, and R. F. Surdick. Quæstiones Entom. 14.
- OBERNDORFER, R.Y. & K. W. STEWART. (1977). The life cycle of Hydropsyche crosbyi (Plecoptera: Perlodidae). Ct Basin Nat. 37: 260-273.
- O'CONNOR, J. P. (1978). The stonefly Capnia atra Norton (Plecoptera, Capniidae) confirmed as an Irish species. Entomol. Gazz. 29: 156-158.
- OKAZAWA, T. (1974). Aquatic insect survey of the River Knunai, with special reference to the rocky chute bed fauna (includes Plecoptera). Kontyu (Tokyo) 43: 497-512.
- OLIVE, J. H. (1976). Chemical-physical and biological assessment of water quality in the Cuyahoga River (1973-74) (includes Plecoptera). Ohio J. Sci. 76: 5-15.
- OLIVE, J. H. & K. R. SMITH. (1975). Renthic macroinvertebrates as indexes of water quality in the Scioto River Basin, Ohio (includes Plecoptera). Bull. Ohio Biol. Surv. 5: 1-24.

- PARKER, N. C. & B. A. SIMCO. (1975). Activity patterns, feeding and behavior of the pirateperch, Aphredoderus sayanus (Plecoptera as food). *Copeia* 1976: 572-574.
- PECKARSKY, B. L. (1979). Biological interactions as determinants of distributions of benthic invertebrates within the substratum of stony streams. *Limnol. Oceanogr. (Ann Arbor)* 24: 59-68.
- PENNAK, R. W. (1977). Trophic variables in Rocky Mountain trout streams. *Arch. Hydrobiol. (Stuttgart)* 80: 253-285.
- RABENI, C. F. & G. W. MINSHAW. (1977). Factors affecting micro-distribution of stream benthic insects. *Oikos (Copenhagen)* 29: 33-43.
- RAVIZZA, C. (1976). Faunistic and taxonomic notes on vernal Plecoptera in Sardinia. *Boll. Soc. Sarda. Sci. Nat.* 15: 149-160. (Ital. with Engl. summ.).
- RAVIZZA, C. (1977). Notes on Protonemura elisabethae Rev., with description of the full-grown nymph (Plecoptera, Nemouridae). *Bull. Soc. Entomol. Ital.* 109: 75-77. (Ital. with Engl. summ.).
- RAVIZZA, C. (1978). Una specie nuova di Plecottero dei monti del biellese: Leuctra brevipennis n. sp. (Plecoptera Leuctridae). Estratto da *REDIA* (Florenz) LXI: 251-257.
- RAVIZZA, C. & E. RAVIZZA DEMATTEIS. (1977a). High Tanaro Valley (Ligurian Alps) stonefly fauna (Plecoptera). *Mem. Soc. Entomol. Ital. (Genova)* 55: 183-215. (Ital. with Engl. summ.).
- RAVIZZA, C. & E. RAVIZZA DEMATTEIS. (1977b). The full-grown nymph of Protonemura ausonia (Cons.) and P. Robusta Berth. *Att. Soc. Ital. Sci. nat. Museo. Civ. Stor. nat. Milano* 118: 353-359.
- RAVIZZA, C. & E. R. DEMATTEIS. (1978). Mountain brooklet stonefly fauna of northern Apennine (Plecoptera). *Atti. Soc. Ital. Sci. nat. Museo. civ. Stor. nat. Milano* 119: 229-242.
- RAYMOND, S. 1979. As I See It: Anglers with fine mesh nets are hauling in Odonata and Plecoptera. *Sports Illus.* VIII: 8-3.

- REICE, S. R. (1977). The role of animal associations and current velocity in sediment - specific leaf litter decomposition. *Oikos* (Copenhagen) 29: 357-365.
- REICE, S. R. (1978). Role of detritivore selectivity in species-specific litter decomposition in a woodland stream. *Verh. Internat. Verin. Limnol.* (Stuttgart) 20: 1396-1400.
- RECKER, W. E. (1976). Origin of stonefly names proposed by Ricker and collaborators. *Perla* 2: 5-19.
- RIEK, E. F. (1976a). A new collection of insects from the Upper Triassic of South Africa. *Ann. Natal. Mus.* 22: 791-820. (Includes Plecoptera).
- RIEK, E. F. (1976b). Fossil insects from the Middle Ecca (Lower Permian) of southern Africa. *Paleontol. Afr.* 19: 145-148. (Paraplecoptera).
- RUPPRECHT, R. (1978). Proof of drumming - signals in a European species of the stonefly family Leuctridae (Plecoptera). *Entomol. Ger.* (Stuttgart) 3: 333-336. (Ger. with Engl. summ.).
- SALMAN, S. A. (1979). Etude écologique d'un ruisseau de plaine "Le Fontanil". Manuskript. Thèse de 1 - Université de Grenoble.
- SALTVEIT, S. J. (1978). The small nymphs of Diura nansenii (Kempny) (Plecoptera). *Ent. Scand.* (Lund) 9: 297-298.
- SALTVEIT, S. J. (1979). Field and laboratory studies on the life history of three species of Amphinemura. *Gewässer und Abwasser* 64: 30-31.
- SANDERS, H. O. & O. B. COPE. (1968). The relative toxicities of several pesticides to nauplii of three species of stoneflies. *Limnol. Oceanogr.* (Lawrence, Kansas) 13: 112-117.
- SANDROCK, F. (1978). A comparison of the insects emerging from two brooks in the region of Schiltz: Breitenbach and Rohrwiesenbach, West Germany 1970-1971. *Arch. Hydrobiol. Suppl.* 54: 328-408.

- SARAI, D. S. (1976). Total and fecal coliform bacteria in some aquatic and other insects (includes Acroneura evoluta Klap.). Environ. Entomol. 5: 365-367.
- SAVAGE, N. L. & F. W. RABE. (1973). The effects of mine and domestic wastes on macroinvertebrate community structure in the Coeur d' Alene River (includes Plecoptera). Northwest Sci. 47: 159-168.
- SCHOENEMUND, E. (1911). Über die hermaphrodite sexualanlage der Männchen von Perla marginata Panz. Sber. Medizin. - Natwiss. Ges. Münster 1911: 1-8. Note: quoted from reprint, overlooked in all catalogues; despite identity of title not the same as paper quoted "1912b" by ILLIES 1966.
- SCHIUSTER, K. (1978). Das zoobenthos der oberen Mur. Decheniana (Bonn) 131: 141-146.
- SHELDON, A. L. & M. W. OSWOOD. (1977). Blackfly (Diptera: Simuliidae) abundance in a lake outlet: test of a predictive model. Hydrobiologia (The Hague) 56: 113-120. (Plec. discussed).
- SHIORT, R. A. & P. E. MASLIN. (1977). Processing of leaf litter by a stream detritivore: effect on nutrient availability to collectors. Ecology 58: 935-938.
- SIEGRIED, C. A. & A. W. KNIGHT. (1976). Prey selection by a setipalpian stonefly nymph, Acroneuria (Calineuria) californica Banks (Plecoptera: Perlidae). Ecology 57: 603-608.
- SIEGFRIED, C. A. & A. W. KNIGHT. (1978). Aspects of the life history and growth of Acroneuria (Calineuria) californica in a Sierra foothill stream. Ann. Ent. Soc. Am. 71: 149-154.
- SIEGFRIED, C. A., P. J. SHIELIAN & A. W. KNIGHT. (1977). The adults of Oroperla barbara (Needham) (Plecoptera: Perlodidae). Pan-Pac. Entomol. (San Francisco) 53: 126-128.
- SIMMONS, G. M. & J. R. VOSHELL, Jr. (1978). Pre- and post-impoundment benthic macroinvertebrate communities of the North Anna River. Current perspectives on river-reservoir ecosystems, North American Bentholog. Soc. ed. 45-61.

- SINGH, R. K. & P. D. RANE. (1976). A new record of stoneflies (Plecoptera: Perlidae) from Madhya Pradesh, India. *Newsl. Zool. Surv. India (Calcutta)* 2: 159.
- SIVEC, I. (1979). Notes on the drift of stoneflies (Plecoptera) in subterranean environment of the sinking-river Pivka (Slovenija, Yugoslavia). *Gewasser und Abwasser* 64: 39-44.
- SLADECEK, V. (1973). System of water quality from the biological point of view. *Arch. Hydrobiol., Beih. Ergebn. Limnol. (Stuttgart)* 7: I-IV, 1-218.
- SILFER, E. H. (1979). Sense organs on the antennal flagellum of a stonefly (Plecoptera, Insecta). *Journal of Morph.* 160: 1-6.
- SNELLEN, R. K. (1979). The life cycle of Perlesta placida (HAGEN). *Gewasser und Abwasser* 64: 29-30.
- SNELLEN, R. K. & K. W. STEWART. (1979). The life cycle and drumming behavior of Zealeuctra classseni (Frison) and Zealeuctra hitei Ricker and Ross (Plecoptera: Leutridae) in Texas, USA. *Aquatic Insects (Lisse)* 1: 65-89.
- SOLON, B. M. & K. W. STEWART. (1972). Dispersal of algae and protozoa via the alimentary tracts of selected aquatic insects. *Environ. Entomol.* 1: 309-314.
- STANFORD, J.E. (1979). Taxonomy and ecology of Rocky Mountain Isocapnia, with special reference to the significance of intraspecific dwarfing. *Gewasser und Abwasser* 64: 76-77.
- STARK, D. P. & R. W. BAUMANN. (1978). New species of Nearctic Neoperla (Plecoptera: Perlidae), with notes on the genus. *Ct Basin Nat.* 38: 97-114.
- STARK, D. P. & A. R. GAUPIN. (1979). The stoneflies (Plecoptera) of Florida. *Trans. Am. Ent. Soc. (Washington)* 104: 391-433.

- STAUFFER, J. R., H. A. BEILES, J. W. COX, K. L. DICKSON, & D. E. SIMONET. (1974-76). Colonization of macrobenthic communities on artificial substrates. *Rev. Biol. (Lisb.)* 10: 49-62.
- STEINMANN, H. (1967). The flying period of Hungarian Plecoptera. *Fol. Entomol. Hung. (Budapest) Stern. Nov.* 20: 197-202.
- STEVENSOM, E. (1977). Growth rates of the stonefly *Amphinemura sulcicollis* Stephens, in the upper River Wear. *Freshwat. Biol. (Oxford)* 7: 455-460.
- STEWART, K. W. & R. L. FULLER. (1979). Food habits of the stoneflies of Gunnison and Dolores Rivers, Colorado. *Gewasser und Abwasser* 64: 45-46.
- STONEBURNER, D. L. (1977). Preliminary observations of the aquatic insects of the Smoky Mountains: altitudinal zonation in the spring. *Hydrobiologia (The Hague)* 56: 137-144.
- SURDICK, R. F. (1979). Preliminary notes on the systematics of the Nearctic Chloroperlidae. *Gewasser und Abwasser* 64: 59-60.
- SURDICK, R. F. & A. R. GAUFIN. (1978). Environmental requirements and pollution tolerance of Plecoptera. U.S.E.P.A. Environmental Monitoring and Support Laboratory. EPA-600/4-78-062. 417 pp.
- SZCZYTOK, S. W. & K. W. STEWART. (1977). The stoneflies (Plecoptera) of Texas. *Trans. Am. Ent. Soc.* 103: 327-378.
- SZCZYTOK, S. W. & K. W. STEWART. (1978). *Isoperla bilineata*: Designation of a Neotype and Allotype, and further descriptions of egg and nymph. *Ann. Ent. Soc. Am.* 71: 212-217.
- SZCZYTOK, S. W. & K. W. STEWART. (1979a). Studies of western Nearctic *Isoperla*. *Gewasser und Abwasser* 64: 73-74.
- SZCZYTOK, S. W. & K. W. STEWART. (1979b). The genus *Isoperla* (Plecoptera) of western North America; holomorphology and systematics, and a new stonefly genus *Cascadoperla*. *Am. Ent. Soc.* 32: 1-120.

- THORUP, J. & C. LINDEGAARD. (1977). Studies on Danish springs. *Folia Limnol. Scand.* 17: 7-15.
- TIMMS, B. V. (1978). The benthos of seven lakes in Tasmania. *Arch. Hydrobiol. (Stuttgart)* 81: 422-444.
- TKAC, M. A. & B. A. FOOTE. (1978). Annotated list of stoneflies (Plecoptera) from Stebbing Gulch in northeastern Ohio. *Great Lakes Entomol.* 11: 139-142.
- TORIAS, W. (1976). Kocherfliegen und Steinfliegen einiger Gewässer in Sor Varanger (Nord-Norwegen) (Trichoptera, Plecoptera). V. Limnephilus minutulus (Banks 1907). *Entomol. Z.* 86: 121-125.
- TOMKIEWICZ, S. M., Jr. & W. A. DUNSON. (1977). Aquatic insect diversity and biomass in a stream marginally polluted by acid strip mine drainage. *Water Res. (Oxford)* 11: 397-402.
- TOWNS, D. R. (1979). Composition and zonation of benthic invertebrate communities in a New Zealand Kauri forest stream. *Freshwat. Biol.* (Oxford) 9: 251-262.
- USSING, H. (1918). Insektilveti of ved Guldenaens Delta og Randers Fjord. In: JOHANSEN, A. C., Randers Fjords Naturhistorie Kobenhaven: 351-392. Nach KAISER 1977: darin offenbar erste Beschreibung der Larve von Isoptena.
- VEIGL, E. (1979). Autecological studies on the development of Plecoptera in a Flysch-stream warming up in summer. *Gewässer und Abwasser* 64: 31-34.
- VERNEAUX, J. & M. REZZOUK. (1974). Les structures d'un grand cours d'eau à Salmonides: la Loue (Massif du Jura). Essai typologique et problème de relations entre espèces et milieu. *Ann. Limnol.* (Paris) 10: 131-162.
- WALTON, O. P. Jr., S.R. REICE & R. W. ANDREWS. (1977). The effects of density, sediment particle size and velocity on drift of Acroneurius abnormis (Plecoptera). *Oikos (Copenhagen)* 28: 291-298.

- WARD, J. V. & R. A. SHORT. (1978). Macroinvertebrate community structure of four special lotic habitats in Colorado, U.S.A. Verh. Internat. Verein. Limnol. (Stuttgart) 20: 1382-1387.
- WHITE, T. R., P. H. CARLSON & R. C. FOX. (1979). Emergence patterns of fall and winter stoneflies (Plecoptera: Filipalpia) in north-western South Carolina. Proc. Entomol. Soc. Wash. 81: 379-390.
- WHYTTON DA TERRA, L. S. (1979). Notes on the Portuguese Plecoptera. Gewasser und Abwasser 64: 60-68.
- WICHARD, W. (1979). Chloride cells of Plecoptera larvae. Gewasser und Abwasser 64: 45.
- WIEMERS, W. (1978). Die invertebratenfauna der Fließgewässer im Solinger Raum. Decheriana (Bonn) 131: 172-182.
- WILHJM, J., H. NAMMINGA, & C. FARRARIS. (1978). Species composition and diversity of benthic macroinvertebrates in Greasy Creek, Red Rock Creek and the Arkansas River. Am. Midl. Nat. 39: 444-453.
- WILLIAMS, D. D. & J. H. MUNDLE. (1978). Substrate size selection by stream invertebrates and the influence of sand. Limnol. Oceanogr. (Ann Arbor) 23: 1030-1033.
- WINTERBOURN, M. J. (1978). The macro invertebrate fauna of a New Zealand forest stream. N.Z. J. Zool. 5: 157-170.
- ZHILTZOVA, L. A. (1977a). New species of stoneflies (Plecoptera) from Primorye Territory. Trudi Zool. Inst. A. N. SSSR (Leningrad) 70: 3-9.
- ZHILTZOVA, L. A. (1977b). The fauna and ecology of stoneflies (Plecoptera) of the Ukrainian Carpathian Mountains. Entomol. Obozr. (Leningrad) 56: 43-52. (Russ. with Engl. summ.).
- ZHILTZOVA, L. A. (1977c). Materials on the stonefly fauna (Insecta, Plecoptera) of the upper parts of drainages of streams in the areas of Ussuri and Primorye. Trudi biol. potschvennogo Inst. (Vladivostok) 46: 17-27. (Russ.).

- ZHILITZOVA, L. A. (1978a). Addition to stonefly fauna of the Nemouridae family (Insecta, Plecoptera) from Central Asia: 2. Vestn. Zool. 1: 35-42. (Russ. with Engl. summ.).
- ZHILITZOVA, L. A. (1978b). New species of stoneflies of the family Chloroperlidae (Plecoptera) from the Soviet Far-East USSR. Entomol. Obozr. 57: 543-548.
- ZHILITZOVA, L. A. (1979). Plecoptera. pp. 438-439 In: MORDUKHAI-BOLTOVSKI, (ed.): The River Volga and its life. Monographiae Biologicae 33, XVI + 473 pp.
- ZHILITZOVA, L. A. & I. M. LEVANIDOVA. (1978). New species of stoneflies (Plecoptera) from the Far East. Trudi Zool Inst. A. N. SSSR (Leningrad) 71: 3-29. (Russ.).
- ZHILITZOVA, L. A. & Y. I. ZAPEKINA-DULKETT. (1977a). The first find of a male of the genus *Baikaloperla* (Plecoptera: Capniidae). Zool. Zh. (Moscow) 56: 307-308. (In Russ. with Engl. summ.).
- ZHILITZOVA, L. A. & Y. I. ZAPEKINA-DULKETT. (1977b). New species of stoneflies (Insecta, Plecoptera) from South Primorje. Freshwater fauna of the reserve "Kedrovaya Pad", Proceedings of the Institute of Biology and Pedology (Vladivostok) 45: 50-52.
- ZHILITZOVA, L. A. & Y. I. ZAPEKINA-DULKETT. (1977c). Larvae of the Baikal stonefly *Baikaloperla kozhovi* Znp. - Dulk. et Zhiltz. (Plecoptera, Capniidae). A species endemic to the Baikal region. Entomol. Obozr (Leningrad). 56: 781-784. (Russ.).
- ZWICK, P. (1977). Plecoptera emerging from two streams at Lunz, 1972-1974. Arch. Hydrobiol. (Stuttgart) 80: 458-505. (Ger. with Engl. summ.).
- ZWICK, P. (1978). Steinfliegen (Insecta, Plecoptera) aus Griechenland und benachbarten Landern. Mitt. schweiz. ent. Ges. (Zürich) 51: 21-38, 213-239.
- ZWICK, P. (1979a). Revision of the stonefly family Eusthenidae (Plecoptera), with emphasis on the fauna of the Australian Region. Aquatic Insects (Liess) 1: 17-50.

ZWICK, P. (1979b). Plecoptera from Greece. *Ceyerat und Abwasser* 64: 69.